



African Journal of Biological Sciences



Advances In Tissue Culture And Somatic Embryogenesis In Fruit Crops.

Jagruti Singh¹, Susmita Das², Dhrubajyoti Banerjee³, Vishakha Saini⁴, Priya Singh⁵, Jatinder Singh^{6*}

^{1,2,3,4,6} Department of Horticulture, School of Agriculture, Lovely Professional University, Phagwara, Punjab (India),

⁵ Department of Computer Science and Engineering, School of Computer Science and Engineering, Lovely Professional University, Phagwara, Punjab (India)

*Corresponding author: Jatinder Singh

*Department of Horticulture, School of Agriculture, Lovely Professional University, Phagwara, Punjab (India)

Email:-jatinder.19305@lpu.co.in

Article History

Volume 6, Issue 3, 2024

Received: 22 Feb 2024

Accepted : 24 Mar 2024

Doi: 10.48047/AFJBS.6.3.2024.268–283

Abstract: –

In modern era, traditional propagation techniques are overpowered by the advanced in vitro culture, that offers the highest commercial value of the plants, due to its rapid generation of clonal plants. Additionally, it is very helpful in supporting the preservation of germplasm, the creation and upkeep of disease-free planting material, and additions to standard plant advancement techniques such as plant genetic engineering. Micropropagation approaches are frequently based on tissue culture techniques. The main objectives of these techniques are to increase the effectiveness and efficiency of plant regeneration. The maximum and minimum use of a few fundamental concepts related to plant tissue culture in vitro, particularly micropropagation, have been discussed in this article. These concepts include types of tissue culture and recent advances for general studies of micropropagation, callus induction, shoot and root induction, and somatic embryogenesis (SE). Additionally, somatic embryogenesis is a significant breakthrough in plant vegetative propagation technology, with implications for high-value horticulture crops and tree breeding. Within a limited amount of time and resources, the technique depicts the mass proliferation of new and elite types. Conventional breeding techniques multiply slowly and rely on natural processes, but somatic embryogenesis may help better in terms of quick mass production of plantlets in a limited time. There are several practical and economic uses of the rapidly evolving somatic embryogenesis techniques, especially in the area of in vitro clonal micropropagation. The low frequency of somatic seedling recovery and the poor germination of its embryos have hampered the application of somatic embryogenesis in fruit crop improvement. In this review, the advances achieved till date, have been elaborated more efficiently, which can effectively utilized for achieving future perspectives.

Key words--- Somatic embryogenesis, Tissue culture,

A. Introduction: –

Micropropagation refers to the aseptic in-vitro cultivation of cells, tissues, organs, and their constituent parts under certain chemical and physical conditions (Gupta et al., 2020). Considered to be the plant regeneration method with the least genetic instability and soma clonal variation is micropropagation through the growth of axillary buds (Encina et al., 2023). Plant Tissue Culture is important for agriculture and plant breeding since it enhances crop production through micropropagation, hybridization, synthetic seed production, haploid culture, and pathogen

eradication (Gulzar et al., 2020). In order to enable micropropagation from altered passion fruit cells, regeneration systems have been constructed from the shoot apex, leaf discs, nodal, and hypocotyl segments, as well as the roots (da Silva et al., 2021). For the study of the morphophysiological, biochemical, and molecular processes that take place during the development of higher plants, somatic embryogenesis makes it simple to manipulate and control the culture conditions (Gerdakaneh et al., 2021). Protoplasts are the parts of plant cells that are enclosed by their cell walls. By physically or by an enzymatic digesting process, the cell membrane can be removed, allowing for plasmolysis and separation. The preservation of this precious biological resource would greatly benefit from the development of a fruitful propagation mechanism. (Cobo et al., 2018). For the sustainable and ongoing synthesis of bioactive chemicals from this threatened species, callus culture is an option (Koufan et al., 2020). A thorough investigation of the genetic and epigenetic factors causing callus induction is warranted, as well as the general use and understanding of molecular mechanisms (Iqbal et al., 2019). Somatic hybridization is the in vitro fusing of plant protoplasts produced either from somatic cells of the same plant or from two genetically distinct plants (Chakraborty et al., 2016). Indirect and Direct SE relate, respectively, to the presence or absence of a callus formation phase (Joshi et al., 2013). Since the first reports on carrot in 1958, somatic embryogenesis (SE) has been reported in various plant species (Solís-Ramos et al., 2012). Somatic Embryogenesis (SE) can arise in vitro either directly from the explant without any intermediary callus stage (direct SE) or indirectly from the callus (indirect SE). In contrast to primary SE created from explant cells, somatic embryogenesis can also be directly triggered or by callus in the culture of somatic embryos. This process is referred to as secondary SE (Solís-Ramos et al., 2012). When mass producing synchronized developmental stages of somatic embryos, somatic embryogenesis (SE) is more advantageous than zygotic embryogenesis since it is simpler to monitor and manage the cultures (Botini et al., 2021). One of the most important components in the induction of somatic embryogenesis in plants is growth regulator, which has been detected in around 80% of somatic embryogenesis (SE) induction techniques. Somatic embryogenesis is caused by variations in the concentration of known growth regulators, specifically auxins and cytokinin, in the plant tissue culture media (Gerdakaneh et al., 2021). Axillary shoot culture and plantlet recovery after shoot regeneration from somatic embryos are two other methods that have been used. After genetic transformation operations in this species, regeneration, and plant recovery frequently take place (Botini et al., 2021). The greatest potential for totipotent growth is seen in in vitro tissue culture, where a wider range of explants may be stimulated to undergo embryogenesis, including haploid cells of the male and female gametophyte (gametophytic embryogenesis) (Horstman et al., 2017). The ability to create plant material without regard to the time of year or the plant's growth cycle is made possible by tissue culture. Additionally, by altering the environment, the genes, and the chemicals in a tissue culture system, it is possible to increase the concentration and purity of therapeutic (Menbari et al., 2021). The development of embryogenic cell suspension cultures requires the selection of large quantities of high-quality embryogenic callus; embryogenic calli and transparent pro embryos should make up the initiating tissues, whereas compact calli, cotyledonary stage embryos, and meristematic globules should be avoided (Kong et al., 2020). Plant protoplasts are living, naked cells surrounded by a plasma membrane that, in the right culture conditions, can lead to the regeneration of plants (Xiangmin et al., 2019). As a source of protoclonal variation, also known as soma clonal variation, protoplast culture could be of great benefit in enhancing the existing germplasm by producing variations that can be applied in breeding programs (Barceló et al., 2019). For many crops, in vitro anther culture is the preferred method of regeneration because to the high responsiveness of explants, the ability to stimulate somatic embryogenesis, and the

potential to regenerate haploid plants from microspores. Many variables, including the anthers physiological state, stage of differentiation, endogenous and external hormonal signalling, and culture medium composition, affect the anthers reactivity in in vitro situations (dos Santos et al., 2021). Meristem culture is a popular technique for removing viruses from horticultural plants. Meristems are thought to be virus-free because of the uneven distribution of viruses in plants. Determining out the meristem tip's size might be helpful in improving the effectiveness of viral eradication. (Vivek et al., 2018).

Different types of Plant Tissue Culture

Tissue culture can be divided into various types as described in Table no 1 and it depends upon the plant part which is used explant for preparing the culture.

Table 1. Types of Tissue Culture

Sr.no	Types of tissue culture based on the part of plant used as explant	Reference	Sr No	Types of tissue culture based on the part of plant used as explant	Reference
1	Apical meristem culture	(Gaikwad et al., 2020)	7	Root & Shoot tip culture	(Gupta et al., 2020)
2	Axillary bud culture	(Gaikwad et al., 2020)	8	Flower organ culture	(Gupta et al., 2020)
3	Callus culture	(Gaikwad et al., 2020)	9	Fruit organ culture	(Gupta et al., 2020)
4	Cell Suspension culture	(Gaikwad et al., 2020)	10	Anther culture	(Gupta et al., 2020)
5	Protoplast culture	(Gaikwad et al., 2020)	11	Somatic Embryogenesis	(Singh et al., 2019)
6	Embryo culture	(Gaikwad et al., 2020)			

1. Apical meristem culture

The most promising method for removing viruses from plant tissue culture as well as for replicating a chosen plant material true to its type and exhibiting the same agronomic properties is plant regeneration through the apical meristem. Shoot tip explants with a single apical meristem responded to regeneration less well than those with two axillary meristems (Sahraroo et al., 2019). The shoot apical meristem (SAM) and root apical meristem (RAM) are formed at the highest point of embryogenesis, while the lowest point is situated at either the shoot top or the root bottom (Xue et al., 2020). Larger meristems could not completely eliminate the viruses under study (Vivek et al., 2018). The details of in- vitro studies of different fruit crop by using apical meristem are given in table no 2.

Table 2. In-vitro studies of different fruit crop by using apical meristem

Species cultivar	Explant used	Culture medium, PGR,	Culture condition	Result	References
Purple passion fruit	Apical meristem	MS medium, BA, NAA, sucrose, IBA		Root formation was satisfactory with IBA @ 0.4 and 0.6 mg/l	Prammanee et al., 2011
Apple	Apical meristem	MS medium supplemented with BA (1.0 mg/l), IBA (0.05 mg/l) and GA3 (0.1 mg/l)	Temp 25 ± 2°C. It was kept at 16-hour light and 8-hour dark photoperiod.	Mostly explants are established, and more number of meristems proliferated with BA (0.5 mg/l), IBA (0.08 mg/l), and GA3 concentration	Vivek et al., 2018

Fig	Apical meristem	MS medium, BA NAA, and IBA was used for shoot regenerating.	pH 5.8 culture was kept at $25 \pm 2^\circ\text{C}$ under 16 h photoperiod,	More numbers of rooted micro-shoots were observed with 1.5mg/l IBA	Sahararo et al., 2019
Wild Guava	Apical meristem	JADS culture medium, myo-inositol, PVP, sucrose, agar	pH 5.7, autoclaving at 121°C for 20 min	The culture media containing 2.22 μM BAP exhibited the highest responsiveness and the maximum number of shoots per explant.	dos Santos et al., 2021
Strawberry	Apical meristem	MS basal medium enhanced with 0.2 mg/l IAA and 1 mg/l BA	Incubated under 16/8 h light/dark cycle at $25 \pm 2^\circ\text{C}$ by cool white fluorescent lights	The medium added with 500 mg/l polyvinylpyrrolidone (PVP) produced the maximum results.	Hemmati Asl, S., & Dorani, E. (2023)

2. Axillary bud culture.

One of the various methods for cultivating plants in vitro is xylem bud culture. One of the main issues with in vitro culture systems is the occurrence of somaclonal variation among subclones of a single parental line. Numerous characteristics can be associated with this variant, such as transposon and retrotransposon activation, sequence modifications, cytological abnormalities, frequent qualitative and quantitative phenotypic alterations, and gene silencing and activation. Therefore, somaclonal variation is taken into account at both the genetic and epigenetic levels. The axillary bud culture approach is commonly thought to present the least danger of genetic instability because of meristems' greater resistance to genetic alterations compared to disordered tissues (Ngezahayo F., & Liu, B. 2014). These techniques still have the potential to produce more plant material faster, cheaper, and use less labour. Over four weeks, these in vitro propagation techniques produce, on average, four to six shoots from a single source shoot. Considered to be the in vitro plant regeneration method with the least genetic instability and soma clonal variation is micropropagation through the growth of axillary buds. Papaya species have benefited greatly from the widespread use of tissue culture-based micropropagation, which provides advantages over traditional propagation techniques. Various strategies have been used, including axillary shoot cultivation. Considered to be the in vitro plant regeneration method with the least genetic instability and soma clonal variation is micropropagation through the growth of axillary buds (Encina et al., 2023). The details of in-vitro studies of different fruit crops using axillary bud are given in table no 3.

Table 3. In-vitro studies of different fruit crop by using Axillary bud to develop culture

Species, Cultivar	Explant used	Culture medium, PGR,	Culture condition	Result	References
Apple	Axillary bud	MS medium, BA and Kn, sucrose, agar, vitamins, NAA	Minimum and maximum temperatures from 6°C to 12°C and 17°C to 25°C	Maximum calli formed at medium supplemented with 2g/l NAA in combination with 2 mg/l cytokinin and higher root length were recorded at 0.25 mg-L of IBA.	Itana et al., 2022
Papaya cv. Solo	Axillary bud	MS medium supplemented with NAA, IBA, and adenine hemisulphate	Cultures were incubated at $25 \pm 1^\circ\text{C}$ under a 16 h & pH 5.7	The rooting medium and the substrate to MS liquid plus 1 mg L ⁻¹ IBA in vermiculite achieved best acclimatization.	Encina et al., 2023

Caucasian apple	Axillary bud	MS medium, glycine, pyridoxin · HCL, thiamine · HCL, nicotinic acid, BA, GA ₃ , IAA, sucrose, plant agar	16-hour photoperiod, 45% relative moisture, and 25 ± 1°C temperature.	The results showed that a rooting percentage of 77.8% was obtained on half strength, LS medium in combination with 0.9 mg L ⁻¹ IBA	Amirchakhmaghi et al., 2019
Blackberry	Axillary bud	BA, NAA, GA ₃ , Agar	pH 5.8 25°C under 16 h light and 8 h dark cycle.	The cultivar "Chester Thornless" had the maximum rate of proliferation, yielding 9.66 shoots on a medium containing a mixture of 2 mg/l BA and 0.2 mg/l IBA.	KEFAYATI et al., 2019

3. Callus Culture

In order to undergo genetic change, callus induction and subsequent plantlet regeneration are necessary (Ikeuchi et al., 2019). The change of cell fate from the somatic state to pluripotency is necessary for callus induction (Ikeuchi et al., 2016). Numerous explant types, including leaves, mature embryos, stems, and immature embryos, can be used to generate callus; nevertheless, the embryogenic potential of various calli varies greatly (Wu et al., 2020). Cotyledon explants have demonstrated high frequency regeneration as callus induction materials because of their increased meristematic tissue composition, less contamination, and enhanced browning. All things considered, explants of leaves and cotyledons are the main sources of callus induction and ensuing plantlet regeneration (Gao et al., 2023). To study plant genetics and improve plant varieties, callus is a desirable target explant (Cimen et al., 2020). The detailed in-vitro studies of different fruit crop by using apical meristem are given in table no 4

Table 4. *In-vitro* studies of different fruit crop by using callus to develop culture

Species, Cultivar	Explant used	Culture medium, PGR	Culture condition	Result	References
Citrus	Ovules extracted from the immature fruits	MS, vitamins, malt extract, sucrose, and supplemented with 2,4-D, BA, and Kn.	pH 5.8, autoclaving at 121°C for 15 min.	The highest callus initiation using EME (MT basal medium + 0.5 gL ⁻¹ malt extract) + Kn (1.0 mg L ⁻¹) were observed	Cimen et al., (2020).
Argan	Leaf and mature fruit samples of argan	Gallic acid and (DPPH), (BF3), 2,4-D, NAA,(KOH), (NaOH), hexane, and ethanol Sucrose, (IAA) ,(BA) and Agar	pH 5.7 autoclave at 121 °C for 20 min. cultures kept at 25 °C	Combination of 1 mgL ⁻¹ NAA and 1 mgL ⁻¹ 2,4-D resulted in the highest callus formation rate	Koufan et al., 2020
Apple	Calli obtained from fruit	MS salt, thiamine, Myoinositol, Glycine Nicotinic acid Pyridoxine,	25 ± 1°C, and a photoperiod of 16 h light: 8h darkne	It was also discovered that MS medium containing 2,4-D + BA produced the finest callus quality, which was friable and white-yellowish.	MenBARI et al., 2021

			ss and light		
Quince	Fruit pulp was used to establish callus	MS media, GBs, sucrose, BA, NAA,	pH 5.8 Incubated @ 25 ± 2 °C in the dark	The ideal growth, maintenance, and induction conditions for quince callus biomass generation were determined to be B5 medium plus 1.77 µM BA and 5.40 µM NAA.	De Bellis et al., 2022
Apple	Fruit pulp was used to establish callus	MS culture, B ₅ medium, sucrose, BA, NAA, 2,4-D	pH 5.8, Incubated at 25 ± 2 °C in dark	When combined with 2.0 mg/L BA and 2.0 mg/L NAA, MS produced the maximum amount of biomass from GD pulp explants, whereas B ₅ combined with 2.0 mg/L BA and 0.2 mg/L 2,4-D produced the highest amount of biomass from MRM pulp explants.	Verardo et al., 2016
Feijoa	Fruit pulp	Cultures were grown on MS; and B ₅ media, sucrose, BA, NAA	pH 5.8, Incubated in the dark at 25 ± 2 °C.	The combination of B ₅ medium containing 0.89 µM BA and 10.7 µM NAA produced the maximum biomass formation of calluses.	Verardo et al., 2019

4. Protoplast culture

Plant protoplasts were isolated for the first time in 1960 by treating cells with enzymes like cellulase, pectolyase, and hemicelluloses that would dissolve cell walls (Sharma et al., 2014). Single, wall-free cells known as protoplasts can divide, multiply, and differentiate by developing calluses before giving rise to the full plant. A flexible system for conducting functional genetic studies and researching a variety of biological functions, including cell shape, membrane function, and hormone signalling, is the protoplast. Protoplasts can also be a valuable biotechnological tool in breeding operations, particularly for vegetatively propagated species like strawberries that have significant levels of polyploidy and heterozygosity. Leaf mesophyll, embryogenic calli, embryogenic suspension cultures, and nonembryonic calli are only a few examples of diverse tissues from which protoplasts can be recovered. From friable embryogenic ovule-derived calli maintained on a growth-regulator-free medium, the protoplasts can be separated (Cimen et al., 2020). By using protoplasts, it is possible to generate combinations of core genes and/or unique cytoplasmic organelles, with the resulting hybrid cells and asymmetric hybrid cells being one in the main applications of cytoplasmic fusion. The protoplast culture can be very valuable since the variance gained in plants regenerated from protoplasts is far more than the variation obtained in plants from structured growth cultures. As a source of proline variation, also known as soma clonal variation, by enlarging the genetic material already present and producing variants that may be employed in breeding initiatives (Barceló et al., 2019). Each protoplast has the potential to regenerate a new wall and perform recurrent mitosis division to produce daughter cells, which may then be used to regenerate viable plants through tissue culture. This process is dependent on the protoplasts receiving the appropriate chemical and physical stimuli. Many species have protoplast-to-plant systems accessible, and their use has been well documented in the literature. Remarkably, not much has changed in terms of the fundamental protocols for protoplast isolation since they were initially documented. Nonetheless, a significant advancement has been achieved in the quantity of species for which protoplast-to-plant systems are available (Davey et al., 2005).

The detailed in– vitro studies of protoplast culture in different fruit crop is given in table no 5

Table 5. In-vitro studies of protoplast culture of different fruit crop

Species, Cultivar or Rootstock	Explant used Size and sourced	Culture medium, PGR, and Additives	Culture condition	Result	References
Strawberry	Shoots	MS medium supplemented with 10 g l ⁻¹ sucrose, 2 mg l ⁻¹ BA and 0.3% Gelrite for 3–4 subcultures	16 h photoperiod, 25 ± 2 °C	Higher shoot regeneration rates in protoplasts cultured in a medium supplemented with TDZ in comparison with BA, 17% of callus regenerating shoot	Barceló et al., 2019
Apple	Leaf mesophyll	MS medium, NAA, BA, mannitol, pectinase, sucrose, KH ₂ PO ₄ , KNO ₃ , CaCl ₂ ·2H ₂ O: MgSO ₄ ·7H ₂ O, KI, CuSO ₄ ·5H ₂ O	pH: 5.8, Incubation periods for 16 h	In the CPW medium, a 20–hour incubation time with a pore size of 25 µm resulted in a maximum yield of protoplast production.	El-Gioushy et al., 2019
Papaya	Shoots	MS medium, Agar, Sugar, Vitamins, BA, NAA, GA ₃	The cultures were kept at 25±1°C Incubation period 16–hour photoperiod.	2.0 mg/L of IBA was found to have the maximum root induction response (86%) and roots suited for acclimatization.	Waidyaratne et al., 2023

5. Anther Culture.

Both internal and exterior cues have an impact on the embryogenic response of anthers in culture. The genotype, physiological state and growth circumstances of donor plants, gamete developmental stage, pre-treatment of flower buds, media, and incubation conditions, as well as the interactions among these parameters, all have a substantial impact on the anther response to in vitro culture. It is likely that anthers from different species, as well as different cultivars within a species, may require quite diverse circumstances in order to go through an embryogenic pathway and mature. There is no one fixed setting or technique for creating plants from anther cultures. Unlike gynogenesis, which uses female gametes like ovules, androgenogenesis, which uses male gametes such as anthers, microspores, or pollen to generate haploids, is androgenesis. The basic concept is to prevent an immature gametic cell from maturing into a gamete and instead induce it to grow into a haploid plant (Das et al., 2018). Haploid embryos were created using parthenogenesis, pseudogamy, and broad hybridization under in vivo conditions. The haploid embryo must be saved using embryo rescue techniques, cultured further to make haploid, and then chromosome doubling must be carried out to produce doubled haploids. For the creation of haploid cells, androgenesis (anther and microspore culture) and gynogenesis (ovary and ovule) were employed, with androgenesis being the preferred method (Das et al., 2018). Another triploid papaya that was derived was divided into tall, semi-tall, and dwarf trees. The triploid strains and the diploid control did not significantly differ in all morphological characteristics. The fruit-bearing triploid strains parthenocarpically. Each variety of dwarf and semi-dwarf had large yields and good bearing. combined with low stature and the triploid strains have a high parthenocarp fruit output could be used to produce fruit commercially (Rimberia, et al., 2007). In androgenesis via anther culture, male gametophyte cells or their progenitors diverge towards a sporophytic phase, resulting in the formation of haploid embryos and plants (Vasanth et al., 2017). When transplanted into a hormone-free media, embryos taken from another culture can grow roots and transform into plantlets. The detailed studies of anther culture in different fruit crops are given

below in Table 6–

Table 6. In-vitro studies of anther culture of different fruit crop

Species, Cultivar or Rootstock	Explant used Size and sourced	Culture medium, PGR, and Additives	Culture condition	Result	References
Wild passion fruit	Anthers	MS medium, B5 vitamins 0.01% (w/v) myo-inositol, 3% (w/v) sucrose, and 0.28% (w/v) h 4.5 μ M BA and different conc 2,4-D	Temp 27 ± 2 °C Incubated at 16-h photoperiod, pH 5.7 ± 0.1 ,	More numerous only when anthers at the DS2 stage were cultivated with 18.1 μ M 2,4-D and 4.5 μ M BA.	da Silva et al., 2021
Apple	Anther	10 μ M BAP, 0.5 μ M naphthaleneacetic acid (NAA), and 5.0% sucrose	Temp 25°C, Incubation at 16 h a day and 8 hours at night	The rate of shoot formation from anthers showed a high-low order of 'Senshu', 'SD', 'ASP', and 'Tsugaru'	Zhang et al., 2017

6. Somatic Embryogenesis

Somatic Embryogenesis is defined as process in which embryo-like structures are formed from somatic tissue and develop into whole plant (Savita et al., 2015). The survival and development of regenerated plants depend on the cultivation environment and are genotype dependent during the complex process of somatic embryogenesis (Morais-Lino et al., 2016). To regenerate clonal plants and create biotechnological breeding tools like transgenic utilized embryogenic cell suspension cultures, somatic embryogenesis is an alternate technique (Savita et al., 2015). This process yields disease-free plantlets which can be applied for breeding, conservation, sanitation, and micropropagation (Meziane et al., 2017). Organogenesis and somatic embryogenesis have generally been employed for transgenic plant regeneration. While organogenesis is the process by which plants regenerate by producing new organs (shoots and roots) on explants, somatic embryogenesis is the process by which bipolar embryos grow from somatic cells. Somatic embryogenesis seems to have several benefits over organogenesis, including the potential for high rates of multiplication, the capacity to scale up using bioreactors, and the ability to distribute using synthetic seeds. As such, somatic embryogenesis has been emphasized as a promising target for gene transfer (Hota et al., 2019). Micropropagation by somatic embryogenesis acts as a powerful tool for genetic improvement of any plant species because of its single cell origin (Jha et al., 2007). Synchronized generation, maturation, and conversion of embryos into plants are essential for propagation, particularly through somatic embryogenesis. Dedifferentiation, chromatin re-modelling, and gene expression programming in somatic cells are all necessary for the phenomenon of somatic embryogenesis to occur (Ali et al., 2017). A typical somatic embryogenesis technique includes many stages such as callus induction, embryogenic callus multiplication, somatic embryos development, and germination. It seems to be influenced by the interaction of the genotype, explant source, media make-up, plant growth regulators, and explant stage at the start of culture. Nevertheless, because a variety of factors affect how an embryogenic callus differentiates, success with the same cultivar might still vary across years (Vasanth et al., 2017). Citrus plants that were recovered in vitro via somatic embryogenesis (SE) from stigmas and styles were true to type, had a brief juvenile phase, and were free of the primary infectious agents (Meziane et al., 2017). To create somatic embryos from embryogenic calli, solid culture media are typically utilized. These media are also useful for observing the many developmental stages of

somatic embryogenesis in date palm. Solid medium, however, cannot be employed to guarantee widespread dispersion. Due to this, every effort has been made to develop embryogenic cell suspension cultures that have successful plant regeneration and a high morphogenetic potential (Abohatem et al., 2017). Plant growth regulators and carbohydrate sources can be changed in a culture medium to boost the rate of cell proliferation, greatly enhancing the pace of regeneration. Although cytokinin like 6-benzyladenine (BA) and kinetin are thought to be necessary for micropropagation processes like embryogenesis, what is necessary for the best regeneration depends on the genotype (Agisimanto et al., 2019). Modern breeding techniques for bananas must take into account somatic embryogenesis. However, somatic embryogenesis is labor-intensive and extremely genotype-dependent. Only a small number of banana cultivars, including the majority of EAHBs, have so far reacted to somatic embryogenesis. Notably, only a small number of cultivars that are receptive to the process have modest frequencies of somatic embryogenesis. To enhance somatic embryogenesis success rates in cultivars that exhibit poor sensitivity to the process, it is necessary to investigate novel methods for inducing somatic embryogenesis in resistant elite cultivars. There is mounting evidence that certain genes, such as transcription factors, control somatic embryogenesis in bananas. These genes are triggered by specific PGRs and environmental stimuli. Somatic embryogenesis and embryogenic cell suspension cultures developed rapidly, and breakthroughs in 1991 completely transformed the landscape. (Adero et al., 2023). The exudation of phenols at higher levels from plant tissues will remove the darkness and resistant character of plant tissues, which will help with tissue regeneration from in vivo sources, which has been hindered by increasing microbial contamination. Citric acid, ascorbic acid, and polyvinylpyrrolidone (PVP) solution soaking and agitation of plant tissue in nutritive medium significantly reduced the phenolic exudation in the medium and helped to form plant tissue from explants from in vivo sources. Due to the lower production of phenolic compounds in seedlings, tall explant seedlings or greenhouse-grown plants (low light and low temperatures) produced less contamination and phenol (Singh et al., 2019)

Type of Somatic Embryogenesis-- Direct somatic embryogenesis

In direct somatic embryogenesis, embryos are formed directly from the cell without the callus formation (Bhatia et al., 2015).




Fig 1. Steps of Direct Somatic Embryogenesis

Indirect somatic embryogenesis

In indirect somatic embryogenesis the callus is formed from the explant and then from the callus tissue embryo is formed (Bhatia et al., 2015)




Fig 2. Steps of Indirect Somatic Embryogenesis

Table 6. *In-vitro* studies of somatic embryogenesis of different fruit crops

Crop name	Explant used	Culture medium, PGR	Culture Condition	Result	References
Pineapple	Crown leaf and Plantlet leaf	BAP + copper + Picloram on MS medium supplemented with vitamins B ₅	Cultures were kept in the dark at 27±1°C 80% humidity for 4 to 6 weeks.	Medium MS with NAA (0.5 mg/l) + BAP (1mg/l) was highly influenced with 8.8 mature somatic embryos per explant.	Cacai et al., 2021
Strawberry	Leaf blade	MS medium containing 3 mg/L TDZ with 0.25 mg/L 2,4-D	Explants were cultured under a 16/8 h light/dark cycle at 25±2°C.	The maximum number of globular embryos per explant obtained for the cv. of Strawberry.	Gerdakaneh et al., 2021
Passion fruit	Anther	2,4-D, 4.5 µM BA, vitamins B ₅ , myo-inositol, sucrose, and Phytigel	Cultures grew for 60 days at 27±2°C, with 16-hour light, 8 hours dark, and 36 µmol m ⁻² s ⁻¹ irradiance.	Embryogenic calluses and embryos were more only when anthers were cultivated in 18.1 µM 2,4-D and 4.5 µM BA.	da Silva et al., 2021
Kinnow mandarin	Nucellus tissues	MS medium, auxins, cytokinin's and malt extract, BAP	Cultures were kept at 25°C with a photoperiod of 16 hours of light. Light intensity was maintained up to 2500 lux.	Highest Somatic Embryo Induction: Medium: MS, BAP:1 mg/l Highest Somatic Embryo Maturation: Medium: MS, 2,4-D:5mg/l BAP: 1 mg/l	Hussain et al., 2016
Guava	Immature ZE	MS medium enhanced with 500 mgL ⁻¹ malt extract, 6% sucrose, 400 mgL ⁻¹ L-glutamine, and 2.0 mgL ⁻¹ 2, 4-D	Incubate at 25 ± 2 °C in dark for 4–6 weeks for induction. Somatic embryos were exposed to a 16/8 hr. photoperiod for plantlet conversion.	SE induction was achieved using explant on MS medium, 2, 4-D and L-glutamine and sucrose. Maximum no of cotyledonary embryos and matured embryos was achieved using MS medium supplemented with polyethylene glycol and sucrose	Bajpai et al., 2016
Guava	endosperm and immature	Basal salts and vitamins supplemented with sucrose, casein	Cultures were maintained in darkness at 23 ±	Ghosi had the maximum rates of embryogenic callus as well	Ameri et al., 2018

	cotyledon.	hydrolysate CH, PGR's like NAA and BA	1 °C.	as the highest rates of embryo maturation and germination, at 77.06% and 37.66%, respectively.	
Papaya	Immature seeds	MS supplemented with 30 g L ⁻¹ sucrose 20 µM ,2,4-D; and 2.0 g L ⁻¹ Phytigel	pH 5.8. dark at 25 ± 1 °C.	The number of somatic embryos at each developmental stage were evaluated during the maturation phase and each developmental stage	Botini et al., 2021
Kinnow Mandarin	Leaves and Nucellar embryo	MT Basal medium with 0.5mg/l BAP and 0.5mg/l Kinetin	pH-5.7 cultures were kept at 25°C under 16/8 hr light/dark with white fluorescent lights.	Protoplast yield: 5.4 x 10 ⁴ /g FW from nucellar embryos, 17.3 x 10 ⁴ /g FW non-viable protoplasts from leaves.	Kazmi et al., 2022
Pomegranate	Axillary shoot	MS and WPM, CoCl ₂ , Casein hydrolysate, NAA and IBA	Kept at 25 ± 2 °C, 16-hour photoperiod with white fluorescent tubes.	Maximum roots (5.60 ± 0.74) and longest (6.78 ± 0.52 cm) on Woody plant medium + 2000 mg ⁻¹ AC.	Desai et al., 2018
Grape	Nodal segment and leaf disc	MS+2.0mgL ⁻¹ 2,4-D+0.5mgL ⁻¹ Kn+30.0gL ⁻¹ sucrose	Darkness at 25±2°C, 16-hour light/8-hour dark, 2000-lux.	Improved root growth with (MS+2.0mgL ⁻¹ 2,4-D+0.5mgL ⁻¹ BA+30.0gL ⁻¹ sucrose). Longer roots with (MS+0.5mgL ⁻¹ IBA+0.5mgL ⁻¹ Kn+15.0gL ⁻¹ sucrose+7.5gL ⁻¹ agar)	Sharma et al., 2018
Grape	axillary bud	½ MS medium containing 2.5 % sucrose, PGR's	16-h light at 25/20°C, 60% RH, or dark at 25°C	No of embryogenic calli is more when there is higher conc of PGR's and 2,4-D/BAP (2:1)	Maillot et al., 2016
Pineapple	Leaflets from crowns	MS medium, basal salts supplemented with of sucrose, vitaminsB5, glycine, glutamine, casein hydrolysate and MgCl ₂ , BAP	Cultures were kept in the dark at 27±1°C humidity of 80%.	55.25% shoot regeneration from somatic embryos using BAP (3 mg/l) + GA3 (2 mg/l)	Cacaï et al., 2021

Conclusion: –

In the recent period of time, tissue culture techniques herald a revolutionary period in horticulture. The development of these methods has significantly increased plant multiplication's effectiveness and scalability, allowing for the rapid mass production of high-quality planting material. Clonal propagation and genetic modification have emerged as critical areas of focus, protecting premium cultivars and guaranteeing the survival of desirable fruit crop traits. Notably, significant advancements in disease control have been made, with tissue culture providing a dependable way to generate pathogen-free plantlets, to combat various biotic stress. Tissue culture has several applications, as demonstrated by its integration with biotechnological technologies, that have ability to respond against abiotic stresses, and significant in the conservation of uncommon and endangered species. For several fruit crops, inefficient seedling regeneration from in vitro culture method remains an important barrier. Future advancements in micropropagation need careful thought, followed by the use of methods that have been designed to attain extremely effective in vitro regeneration. The early procedures employed stress or growth factor treatments for SE

induction and the process was examined at the histology level. SE can currently be induced by embryo- and meristem-expressed tissue factor (TF), and the other factors implicated in wound healing. Making a thoughtful decision on the utilization of designed methods that are extremely effective in vitro regeneration is crucial. The literature suggests that, it might provide incredibly effective suspension cultures of embryogenic cells from selected calluses lines to go over the present obstacles and produce a speedy technique of clonal propagation. Through the callus phase, somatic embryos can develop either directly or indirectly. genetic metamorphosis through the anther, leaf, shoots, cotyledon, and protoplast. Although transformation has been shown several times, more studies using indirect somatic embryo regeneration are still needed. Future research should focus on enhancing in vitro culture through the use of cell suspension culture technologies and medium additions to increase the induction of somatic embryos. Tissue culture has the potential to improve fruit crops' nutritional characteristics in the future, which would increase global food security.

Reference: –

1. Abohaterem, M. A., Bakil, Y., & Baaziz, M. (2017). Plant regeneration from somatic embryogenic suspension cultures of date palm. *Date palm biotechnology protocols volume I: tissue culture applications*, 203–214.
2. Adero, M., Tripathi, J. N., & Tripathi, L. (2023). Advances in somatic embryogenesis of Banana. *International Journal of Molecular Sciences*, 24(13), 10999.
3. Agisimanto, D., Normah, M. N., & Ibrahim, R. (2019). Rapid somatic embryogenesis of *Citrus reticulata* Blanco cv. Madu in an air-lift bioreactor culture. *Agri vita, Journal of Agricultural Science*, 41(2), 284–294.
4. Ali, M., Mujib, A., Tonk, D., & Zafar, N. (2017). Plant regeneration through somatic embryogenesis and genome size analysis of *Coriandrum sativum* L. *Protoplasma*, 254, 343–352.
5. Ameri, A., Davarynejad, G. H., Moshtaghi, N., & Tehranifar, A. (2018). Polyethylene glycol and chilling overcome Somatic embryogenesis obstacles in *Pyrus communis*. *Scientia horticulturae*, 227, 57–64.
6. Amirchakhmaghi, N., Hosseinpour, B., & Yousefzadeh, H. (2019). Development of a micropropagation protocol for *Malus orientalis* using axillary buds. *In Vitro Cellular & Developmental Biology–Plant*, 55, 625–634.
7. Bajpai, A., Kalim, S., Chandra, R., & Kamle, M. (2016). Recurrent somatic embryogenesis and plantlet regeneration in *Psidium guajava* L. *Brazilian Archives of Biology and Technology*, 59, e16150170.
8. Barceló, M., Wallin, A., Medina, J. J., Gil-Ariza, D. J., López-Casado, G., Juárez, J., ... & Pliego-Alfaro, F. (2019). Isolation and culture of strawberry protoplasts and field evaluation of regenerated plants. *Scientia Horticulturae*, 256, 108552.
9. Botini, N., Almeida, F. A., Cruz, K. Z. C. M., Reis, R. S., Vale, E. M., Garcia, A. B., ... & Silveira, V. (2021). Stage-specific protein regulation during somatic embryo development of *Carica papaya* L. 'Golden'. *Biochimica et Biophysica Acta (BBA)–Proteins and Proteomics*, 1869(2), 140561.
10. Cacaï, G. H. T., Ahokpossi, B. A., Houédjissin, S. S., Houngue, J. A., Badou, B. T., & Ahanhanzo, C. (2021). Plant Regeneration Through Somatic Embryogenesis in Two Cultivars of Pineapple (*Ananas Comosus* L). Preprint from Research Square. <https://doi.org/10.21203/rs.3.rs-542908/v1>
11. Chakraborty S. (2016). Protoplast Culture and Somatic Hybridization, a Promising Frontier of

- Plant Biotechnology. J. Arts, Science & Teaching (JAST), ISSN 2395-4353. 2 (1), pp. 46-54.
12. Cimen, B. (2020). Efficient protoplast isolation from ovule-derived embryogenic callus in *Citrus volkameriana*. Turkish Journal of Agriculture and Forestry, 44(6), 567-576.
 13. Cobo, M. M., Gutiérrez, B., & de Lourdes Torres, M. (2018). Regeneration of mortiño (*Vaccinium floribundum* Kunth) plants through axillary bud culture. In vitro cellular & developmental biology-plant, 54, 112-116.
 14. Davey, M. R., Anthony, P., Power, J. B., & Lowe, K. C. (2005). Plant protoplasts: status and biotechnological perspectives. Biotechnology advances, 23(2), 131-171.
 15. Das, A., Kumar, K., Tribhuvan, K., Das, S. S., & Mishra, M. (2018). Development of haploid and double haploid in fruit crops-A review. International Journal of Current Microbiology and Applied Sciences, 7(5), 2119-2132.
 16. da Silva, M. L., Pinto, D. L. P., de Campos, J. M. S., de Carvalho, I. F., Rocha, D. I., Batista, D. S., & Otoni, W. C. (2021). Repetitive somatic embryogenesis from wild passion fruit (*Passiflora cincinnata* Mast.) anthers. Plant Cell, Tissue and Organ Culture (PCTOC), 146, 635-641.
 17. De Bellis, R., Chiarantini, L., Potenza, L., Gorassini, A., Verardo, G., De Marco, R., ... & Fraternali, D. (2022). High production of secondary metabolites and biological activities of *Cydonia oblonga* Mill. pulp fruit callus. Journal of Functional Foods, 94, 105133.
 18. Desai, P., Patil, G., Dholiya, B., Desai, S., Patel, F., & Narayanan, S. (2018). Development of an efficient micropropagation protocol through axillary shoot proliferation for pomegranate variety 'Bhagwa'. Annals of Agrarian science, 16(4), 444-450.
 19. dos Santos Soares, W., Bruckner, C. H., da Cruz, A. C. F., Rocha, D. I., de Matos, E. M., Koehler, A. D., ... & Otoni, W. C. (2021). Shining light on anther culture, a poorly understood regeneration route in passion fruit (*Passiflora gibertii* NE Brown): histological, hormonal, and phytochemical aspects in vitro Cellular & Developmental Biology-Plant, 57(6), 998-1008.
 20. dos Santos, M. A. C., do Rêgo, M. M., de Queiróz, M. A., da Cruz, A. C. F., da Silva, T. C. R., Pinheiro, M. V. M., ... & Otoni, W. C. (2021). Evaluation of root-to-shoot de novo organogenesis in wild guava species, *Psidium schenckianum* and *P. guineense* (Myrtaceae). Vegetos, 34, 68-76.
 21. Encina, C. L., Granero, M. L., & Regalado, J. J. (2023). In vitro Long-Term Cultures of Papaya (*Carica papaya* L. cv. Solo). Horticulturae, 9(6), 671.
 22. El-Gioushy, S. F. E. E. S., Kareem, A., & Baiea, M. H. M. (2019). Pre-isolation, isolation and regeneration protoplasts from leaf mesophyll of in vivo *Malus domestica* 'Anna'cv. Revista Brasileira de Fruticultura, 41, e-561.
 23. Gaikwad, A. V., Singh, S. K., & Gilhotra, R. (2017). Plant tissue culture-A review. Journal of Pharmaceutical Research & Education, 2(1), 217-220.
 24. Gao, L., Liu, J., Liao, L., Gao, A., Njuguna, B. N., Zhao, C., ... & Han, Y. (2023). Callus induction and adventitious root regeneration of cotyledon explants in peach trees. Horticulturae, 9(8), 850.
 25. Gerdakaneh, M., & Mozaffari, A. A. (2021). Plant regeneration via direct somatic embryogenesis in three strawberry (*Fragaria ananassa* Duch.) cultivars. Agrotechniques in Industrial Crops, 1(3), 103-109.
 26. Gulzar, B., Mujib, A., Malik, M. Q., Mamgain, J., Syeed, R., & Zafar, N. (2020). Plant tissue culture: agriculture and industrial applications. In Transgenic technology Based value addition in plant biotechnology (pp. 25-49). Academic Press.
 27. Gupta, N., Jain, V., Joseph, M. R., & Devi, S. (2020). A review on micropropagation culture method. Asian Journal of Pharmaceutical Research and Development, 8(1), 86-93.
 28. Hemmati Asl, S., & Dorani, E. (2023). Effect of various phenol inhibitors and explant size on

- meristem culture in the Gavieta variety of *Fragaria ananassa*. *Journal of Plant Physiology and Breeding*, 13(2), 1–13.
29. Horstman, A., Bemer, M., & Boutilier, K. (2017). A transcriptional view on somatic embryogenesis. *Regeneration*, 4(4), 201–216.
 30. Hota, D., Kalatippi, A. S., Nayak, J. K., & Karna, A. K. (2020). Genetic transformation of fruit crops: A review. *International Journal of Horticulture and Food Science*; 2(1): 49–51
 31. Hussain, M., Raja, N. I., Iqbal, M., Iftikhar, A., Sadaf, H. M., Sabir, S., ... & Faz, M. N. A. (2016). Plantlets regeneration via somatic embryogenesis from the nucellus tissues of Kinnow mandarin (*Citrus reticulata* L.). *American Journal of Plant Sciences*, 7(6), 798–805.
 32. Ikeuchi, M., Favero, D. S., Sakamoto, Y., Iwase, A., Coleman, D., Rymen, B., & Sugimoto, K. (2019). Molecular mechanisms of plant regeneration. *Annual review of plant biology*, 70, 377–406.
 33. Ikeuchi, M., Ogawa, Y., Iwase, A., & Sugimoto, K. (2016). Plant regeneration: cellular origins and molecular mechanisms. *Development*, 143(9), 1442–1451.
 34. Iqbal, M., Wali, V. K., Bakshi, P., Kour, K., Razdan, V. K., Sinha, B. K., & Sood, K. K. (2019). In vitro propagation of Citrus species through callus induction and regeneration: A review. *Int. J. Curr. Microbiol. App. Sci*, 8(10), 2282–2295.
 35. Itana, D. D., & Habib, Z. Y. In vitro micropropagation of apple (*Malus domestica*) through axillary buds and shoot apices culture. *Int. J. Zool. Appl. Biosci.*, 7(3), 1–7
 36. Jha, T. B., Mukherjee, P., & Datta, M. M. (2007). Somatic embryogenesis in *Jatropha curcas* Linn., an important biofuel plant. *Plant Biotechnology Reports*, 1, 135–140.
 37. Joshi, R., & Kumar, P. (2013). Regulation of somatic embryogenesis in crops: a review. *Agricultural Reviews*, 34(1), 1–20.
 38. Kazmi, S. K., Kayani, H. A., RAZIQ, M., Naz, S., Khan, S., & Kabir, N. (2022). Protoplast isolation and regeneration from leaves and nucellar embryos of kinnow mandarin (*Citrus reticulata* BLANCO). *Pak. J. Bot*, 54(4), 1359–1364.
 39. KEFAYETI, N., Kafkas, E., & ERCİŞLİ, S. (2019). Micropropagation of 'Chester thornless' blackberry cultivar using axillary bud explants. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 47(1).
 40. Kong, E. Y., Biddle, J., Foale, M., & Adkins, S. W. (2020). Cell suspension culture: A potential in vitro culture method for clonal propagation of coconut plantlets via somatic embryogenesis. *Industrial Crops and Products*, 147, 112125.
 41. Koufan, M., Belkoura, I., Mazri, M. A., Amarraque, A., Essatte, A., Elhorri, H., ... & Alaoui, T. (2020). Determination of antioxidant activity, total phenolics and fatty acids in essential oils and other extracts from callus culture, seeds and leaves of *Argania spinosa* (L.) Skeels. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 141, 217–227.
 42. Maillot, P., Deglène-Benbrahim, L., & Walter, B. (2016). Efficient somatic embryogenesis from meristematic explants in grapevine (*Vitis vinifera* L.) cv. Chardonnay: An improved protocol. *Trees*, 30, 1377–1387.
 43. Menbari, A., Bahramnejad, B., Abuzaripoor, M., Shahmansouri, E., & Zarei, M. A. (2021). Establishment of callus and cell suspension cultures of Granny Smith apple fruit and anti tyrosinase activity of their extracts. *Scientia Horticulturae*, 286, 110222.
 44. Meziame, M., Frasher, D., Carra, A., Boudjeniba, M., D'Onghia, A. M., Mercati, F., ... & Carimi, F. (2017). Attempts to eradicate graft-transmissible infections through somatic embryogenesis in *Citrus* ssp. and analysis of genetic stability of regenerated plants. *European journal of plant pathology*, 148, 85–95.
 45. Morais-Lino, L. S., Santos-Serejo, J. A., Amorim, E. P., de Santana, J. R. F., Pasqual, M., & de

- Oliveira e Silva, S. (2016). Somatic embryogenesis, cell suspension, and genetic stability of Banana cultivars. *In Vitro Cellular & Developmental Biology-Plant*, 52, 99–106.
46. Ngezahayo F., & Liu, B. (2014). Axillary bud proliferation approach for plant biodiversity conservation and restoration. *Int. J. Biodivers*, 2014, 1–9.
 47. Prammanee, S., Thumjamras, S., ChiemsomBA, P., & Pipattanawong, N. (2011). Efficient shoot regeneration from direct apical meristem tissue to produce virus-free purple passion fruit plants. *Crop Protection*, 30(11), 1425–1429.
 48. Rimberia, F. K., Adaniya, S., Ishimine, Y., & Etoh, T. (2007). Morphology of papaya plants derived via anther culture. *Scientia Horticulturæ*, 111(3), 213–219.
 49. Sahraroo, A., Zarei, A., & Babalar, M. (2019). In vitro regeneration of the isolated shoot apical meristem of two commercial fig cultivars 'Sabz' and 'Jaami-e-Kan'. *Biocatalysis and Agricultural Biotechnology*, 17, 743–749.
 50. Savita, Pati, P. K., Virk, G. S., & Nagpal, A. (2015). An efficient somatic embryogenesis protocol for Citrus jambhiri and assessment of clonal fidelity of plantlets using RAPD markers. *Journal of Plant Growth Regulation*, 34, 309–319.
 51. Sharma, D. & Tripathi, Manoj & Tiwari, Rajesh & Patel, Ram Prabesh & Baghel, Bindushree & Ahuja, Ashok. (2018). Somatic Embryogenesis and Plantlet Regeneration via Embryogenic suspensions of Grape (*Vitis vinifera* L.). 112–125.
 52. Sharma, D. P. (2014). Plant Tissue Culture in Crop Improvement. In *Proceedings of the 2014 Annual Meeting of the International Plant Propagators Society* 1085 (pp. 273–284).
 53. Singh, G., Sahare, H., & Deep, M. (2019). Recent Trends in Guava Propagation-A Review. *Biosciences Biotechnology Research Asia*, 16(1), 143–154.
 54. Solís-Ramos, L. Y., Andrade-Torres, A., Sáenz-Carbonell, L. A., Oropeza-Salín, C. M., & Castaño de la Serna, E. (2012). Somatic embryogenesis in recalcitrant plants. *Embryogenesis*, 597–618.
 55. Vasanth, K., & Vivier, M. A. (2017). Somatic embryogenesis from immature anther explants: toward the development of an efficient protocol production of grapevine. *Sustainable Agriculture towards Food Security*, 291–305.
 56. Verardo, G., Gorassini, A., Ricci, D., & Fraternal, D. (2017). High triterpenic acids production in callus cultures from fruit pulp of two apple varieties. *Phytochemical Analysis*, 28(1), 5–15.
 57. Verardo, G., Gorassini, A., & Fraternal, D. (2019). New triterpenic acids produced in callus culture from fruit pulp of *Acca sellowiana* (O. Berg) Burret. *Food Research International*, 119, 596–604.
 58. Vivek, M., & Modgil, M. (2018). Elimination of viruses through thermotherapy and meristem culture in apple cultivar 'Oregon Spur-II'. *Virus Disease*, 29, 75–82.
 59. Waidyaratne, S. M., & Samanmalie, L. G. (2022). In vitro Propagation of *Carica papaya* L. Variety 'Horana Papaya Hybrid 01' Using Shoot Tip. *World*, 10(1), 15–9.
 60. Wu, G. Y., Wei, X. L., Wang, X., & Wei, Y. (2020). Induction of somatic embryogenesis in different explants from *Ormosia henryi* Prain. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 142, 229–240.
 61. Xiangmin, S. U. O., Jie, H. A. O., Xinmin, Y. A. N., Xiang, W. A. N. G., Jianzhong, F. E. N. G., & Xueying, L. I. (2019). Protoplast Culture and Its Application in Fruit Breeding. *Agricultural Biotechnology* (2164–4993), 8(3).
 62. Xue, Z., Liu, L., & Zhang, C. (2020). Regulation of shoot apical meristem and axillary meristem development in plants. *International Journal of Molecular Sciences*, 21(8), 2917.
 63. Zhang, C., Sato, S., Tsukuni, T., Sato, M., Okada, H., Yamamoto, T., ... & Komori, S. (2017). Elucidating cultivar differences in plant regeneration ability in an apple anther culture. *The*

Horticulture Journal, 86(1), 1–10